

# OZONE OXIDATIVE POST-CONDITIONING IN ACUTE RENAL FAILURE

José Luis Calunga<sup>1</sup>, Silvia Menéndez<sup>1</sup>, Zullyt Zamora<sup>1</sup>, Nelson Merino<sup>2</sup> and Teresa Montero<sup>3</sup>

<sup>1</sup>Ozone Research Center. PO Box 6414, Havana, Cuba.

E-mail:silviamenendez@infomed.sld.cu

<sup>2</sup>Center for Pharmaceutical Chemistry, Havana, Cuba.

<sup>3</sup>“Luis Díaz Soto” Military Medicine Institute, Havana, Cuba.

## Abstract

The ischaemia/reperfusion process is largely mediated by reactive oxygen species. Taking into account that a transient and controlled administration of ozone is able to upregulate cellular antioxidant enzymes, a morphological, biochemical and functional renal study was performed in rats undergoing a warm renal ischaemia. Rats were divided into four groups. All except the negative controls underwent 60 min´ bilateral renal ischaemia followed by 10 days´ reperfusion. The positive control group received no further treatment. The ozone group received an ozone/oxygen mixture (ozone dose 0.5 mg/kg) immediately after the ischemia and daily for the 10 days´reperfusion; the oxygen group were given the same concentration of oxygen alone (13 mg/kg). Biochemical parameters fructosamine, phospholipase A<sub>2</sub>, catalase, superoxide dismutase and thiobarbituric acid reactive substances were measured, as well as renal plasma flow and glomerular filtration rate. The results demonstrated that, renal plasma flow and glomerular filtration rate decreased significantly in the positive controls and the oxygen group whereas values in the ozone group were similar to those to the negative control group. With respect to the biochemical parameters, ozone maintained a homeostasis redox, with significant increases in catalase and superoxide dismutase activities and similar values for phospholipase A<sub>2</sub> and fructosamine compared with the negative control group. Fewer morphological alterations were seen in kidneys from the ozone group. No advantages were obtained in the positive control and oxygen groups. The protective effect of ozone may be explained by upregulating of the antioxidant defence system and beneficial effects on blood circulation and in oxygen metabolism. Ozone treatment may represent a therapeutic approach for minimizing renal damage after transplantation.

**Key words:** Ozone therapy; reactive oxygen species; reperfusion; superoxide dismutase; warm ischaemia.

## Introduction

Changes that occur during ischaemia affect the cells and provide the basis for increased damage during reperfusion. One of the characteristics of cells submitted to ischaemia-reperfusion (I/R) is that they can recover and have a normal function if the damage is not lethal (no lesion in the tubular basal membrane) or where attempts are made to stop the routes of this damage [1].

Renal vasoconstriction and tubular dysfunction have been clearly defined as the most important mechanisms that reduce the glomerular filtration rate (GFR) after I/R [1]. Decreases in renal blood flow of 40 to 50 % have been reported after a renal ischaemia, in experimental models and in humans. This decrease provokes an increase in shearing force in the endothelium, as well as of neutrophils and reactive oxygen species (ROS). All these effects increase endothelial damage, favouring the maintenance of vasoconstriction and therefore decreasing GFR [2,3].

At present, a protagonist role has been appointed to free radicals, in that increasing prooxidant substances produces an imbalance with antioxidant substances, creating oxidative stress [4,5]. The main oxidative products in biological systems are the ROS which include superoxide anion, hydrogen peroxide, hydroxyl radical, singlet oxygen and hypochlorous acid [6,7]. ROS are implicated in renal damage caused by I/R [3-5]. When ROS are increased and not neutralized with antioxidant substances, they are capable of inducing lipid peroxidation in membranes. They can react with unsaturated fatty acid, provoking structural and functional modifications. In addition, ROS oxidise the lateral chains of amino acid residues and stimulate the formation of protein-protein cross bonds, as well as the oxidation of the protein skeleton resulting in fragmentation.

Some researches have tried to neutralize ROS damage using antioxidant enzymes such as SOD, CAT and glutathione peroxidase, or allopurinol (xanthine oxidase inhibitor) and deferoxamine (inhibitor of the Fenton reaction). They obtained reductions in cell damage and organ dysfunction caused by I/R, proving the role of oxidative stress in the I/R process [7-9]. In a rat model of hepatic I/R, ozone corrected the chronic oxidative stress by upregulating the antioxidant system, achieving an homeostasis redox [10-12].

Repeated non-lethal cell stress has been demonstrated to give protection to a latter severe stress, as occurs in ischaemic, thermal and chemical preconditionings [13-15], as well as in ozone oxidative preconditioning [10-12,16-20].

A comparison between ischaemic and ozone oxidative preconditioning showed no biochemical difference among the antioxidant-prooxidant enzymes measured. Nevertheless, the histological study demonstrated that the protective effect produced by the ozone oxidative preconditioning was superior to that achieved with ischaemic preconditioning [17].

Recent studies have also demonstrated that ischemic postconditioning during reperfusion inhibits myocardial injury (highlighting the myocardial reperfusion phase as a target for cardioprotection) [21,22]. Clinical trials using ozone oxidative postconditioning in cardiopathy and diabetic patients have demonstrated control of oxidative stress, improvements in the antioxidant defence system and satisfactory clinical responses [23,24].

Taking into account that I/R is a process largely mediated by ROS generation and that a controlled and judicious administration of O<sub>3</sub> is able to stimulate the endogenous antioxidant systems [10-12,16-20], and also to enhance blood circulation and oxygen metabolism [6], among other beneficial effects, the aim of this study is to evaluate the efficacy of ozone oxidative postconditioning (applied after the ischaemia) on renal morphology, function and biochemical parameters in rats undergoing a warm renal ischaemia.

## Materials and Methods

### Animals and sample preparation

Forty adult male Wistar rats (250-260 g) were maintained in an air-filtered and temperature-conditioned room (20-22°C), relative humidity 50-52 %. Rats were fed with a standard commercial diet and water *ad libitum*.

Experiments were conducted in accordance with the ethical guidelines established by the Principles of Laboratory Animal Care (NIH publication No. 86-23, revised 1985) and were approved by the Ethical Committee for Animal Experimentation of the National Center for Scientific Research, Havana, Cuba.

Ozone (O<sub>3</sub>) was generated using OZOMED equipment (Ozone Research Center), from medical grade oxygen (O<sub>2</sub>) by means of a silent electric discharge, representing about 3 % of the gas mixture (O<sub>3</sub>+ O<sub>2</sub>). The schedule and O<sub>3</sub> dosing used have been demonstrated to be optimal in previous studies [10-12,16-20].

Under constant sodium pentobarbital anaesthesia, after a renal ischemia of 60 min, we allowed a reperfusion period of ten days (10 d). Heparin (50 U) was administered subcutaneously before induction of ischaemia. Within 10 min of the end of the reperfusion period we measured renal plasma flow (RPF) and glomerular filtration rate (GFR) by means of plasma clearance of p-amino-hippurate (PAH) and inulin, respectively. A constant plasma concentration of both substances was used (2 mg of PAH and 20 mg of inulin in 100 ml of saline solution), given by a continuous perfusion through the left femoral vein at a rate of 0.15 mL/ min, after a loading dose of 0.8 mL of PAH (12 mg/mJ) and 0.8 mL of inulin (2 mg/mJ). For these analyses, blood was withdrawn by intracardiac puncture and urine was collected from the bladder. Rats were then euthanized under deep anaesthesia.

Representative samples of different kidney portions were taken for histopathological studies and preparation of tissue homogenates. Kidney homogenates were obtained using a tissue homogenator Edmund Bulher LBMA at 4°C. The homogenates were prepared in 50 mM KCl/histidine buffer pH 7.4, 1:10 (w/v) and were centrifuged at 8500g at 4°C for 20 min. The supernatants were used for biochemical determinations.

### Treatment schedule and renal ischemia

Animals were divided into four groups of 10 animals. All animals were anaesthetized with sodium pentobarbital (30 mg/kg i.p.) and were given 50 IU of heparin by subcutaneous injection. A laparotomy was then performed. The negative control group underwent sham exposure of the kidneys, followed by resuturing of the abdominal wall, allowing normal renal function for the next 10 days. The other three groups of rats were submitted to bilateral renal ischemia by cross-clamping of both renal arteries for 60 min. Reperfusion thus occurred over the next 10 days, after which the morphological, functional and biochemical renal study was done. The positive control (ischaemia) group received no further treatment. The ozone group were treated with the ozone /oxygen gas mixture administered by rectal insufflation performed with a polyethylene cannula in a volume of 2.5-2.6 mL; the ozone concentration was 50 µg/ml (representing a dose of 0.5 mg/kg weight). Administration was started immediately after 60 min´ renal ischemia and was repeated once daily during the 10-day reperfusion period. The oxygen group were treated in the same way as the ozone group, but insufflating with O<sub>2</sub> (13 mg/kg weight) alone. This group was included to determine the effect of the oxygen in the oxygen/ozone gas mixture.

## **Biochemical determinations**

Levels of PAH and inulin were determined in plasma and urine samples deproteinated by cadmium sulphate. PAH was measured using a photocolometric [25] technique as modified by Smith and Tinkelstein [26]. Inulin was measured by the direct method of resorcinol without alkaline treatment [27]. Plasma creatinine concentrations were measured in deproteinated filtrates using the sodium tungstate method, modified as described by Calunga et al [18].

Kidney homogenates were assayed for total SOD (Cu/Zn and Mn SOD) activity by determining the capacity of the enzyme to inhibit the autoxidation of pyrogallol by 50% [28].

Catalase activity was determined according to the method of Evans and Diplock [29] and the absorbance was measured at 240 nm for 30 s in the spectrophotometer.

A lipid peroxidation assay was used to estimate thiobarbituric acid-reactive substances (TBARS) levels, as described by Ohkawa et al [30] where the absorbance of 3 mL of the colored layer was measured spectrophotometrically at 532 nm, using 1,1,3,3-tetraethoxypropane as standard.

Phospholipase A<sub>2</sub> activity was determined according to a standard procedure [31]; enzymatic activities are expressed as U/g protein.

Protein concentrations were determined by the method of Lowry et al [32], using bovine serum albumin as standard.

Fructosamine was determined by means of a colorimetric procedure [33]. Values represent the difference in units of optical density per g renal tissue.

## **Histological study**

Samples of rat kidneys from the different groups were fixed in neutral 10 % formalin, processed and embedded in paraffin. The histological sections, stained with haematoxylin and eosin, were examined by a pathologist unaware of the treatment schedule.

## **Statistical analysis**

First, the OUTLIERS preliminary test for detection of error values was performed. One-way analysis of variance was then used, followed by the Bartlett-Box homogeneity variance test. In addition, a multiple comparison test was used (Duncan test). Comparisons between two groups were done using the Student's t-test. Results are presented as mean  $\pm$  standard deviation (SD). Differences between groups in the histological study were evaluated with a non parametric test (Fisher's test). Results are presented as means  $\pm$  SD;  $p < 0.05$  was considered significant.

## Results

### Renal function

Table 1 shows the effects of treatment on renal function. RPF shows significant differences between the negative control group and the rest of the groups. Values were lower in the positive control and O<sub>2</sub> groups than in the negative control and the O<sub>3</sub> groups, with significant differences between them. Values were lower in the O<sub>3</sub> group than in the negative control group. Results for GFR showed a similar pattern.

Table 1. Plasma clearance of p-amino-hippurate (PAH), inuline and creatinine

Groups	PAH ( mL/min)	Inuline (mL/min)	Creatinine (µmol/L)
Negative control	1.93 ± 0.006 <sup>a</sup>	0.78 ± 0.08 <sup>a</sup>	83.4 ± 24 <sup>a</sup>
Positive control	0.13 ± 0.003 <sup>b</sup>	0.044 ± 0.010 <sup>b</sup>	177.8 ± 58 <sup>b</sup>
O <sub>3</sub>	0.50 ± 0.11 <sup>d</sup>	0.24 ± 0.07 <sup>d</sup>	90.4 ± 32 <sup>a</sup>
O <sub>2</sub>	0.035 ± 0.0004 <sup>c</sup>	0.013 ± 0.002 <sup>c</sup>	181.9 ± 62 <sup>b</sup>

Values are means ± SD (n = 10 per group). Different letters represent significant differences (p < 0.05) for each biochemical parameter.

With respect to plasma creatinine concentrations, values were higher in the positive control and O<sub>2</sub> groups than in the negative control and O<sub>3</sub> groups. Differences between the positive control and O<sub>2</sub> groups, and between the negative control and O<sub>3</sub> groups, were not significant.

These results demonstrated a deterioration in renal function and that no protection occurs in the positive control and O<sub>2</sub> groups. Thus, oxygen does not favour recovery of renal function. However, ozone oxidative post-conditioning preserved renal function, maintaining values for RPF, GFR and creatinine concentration almost at levels in the negative control group.

### Antioxidant-prooxidant balance and protein oxidation

Table 2 shows the behaviour of the different parameters relating to redox balance in renal tissue. With respect to TBARS, an indicator of lipid peroxidation, the values were significantly higher in the O<sub>2</sub> group than in the other groups. Values in the O<sub>3</sub> and positive control groups were similar, and significantly higher than in the negative control. With regard to SOD, a scavenger of superoxide anions, enzymatic activity was significantly lower in the positive control group than in the other groups. A highly significant increase in SOD was found in the O<sub>3</sub> group. CAT showed similar behaviour to SOD. With respect to fructosamine (an indicator of advanced glycosylation protein products) and the enzyme activity of phospholipase A<sub>2</sub>, significantly higher values were found in the positive control and O<sub>2</sub> groups than in the negative control and O<sub>3</sub> groups. Differences between the negative control and the O<sub>3</sub> groups were not significant.

Ozone oxidative post-conditioning was able to preserve the antioxidant-pro-oxidant balance. Also, protein oxidation was controlled, showing similar behaviour as the negative control group.

Table 2. Biochemical parameters related to redox balance in renal tissue

Parameters	Negative control	Positive control	O <sub>3</sub>	O <sub>2</sub>
CAT (k <sub>15</sub> /g of wet tissue)	5.70 ± 0.30 <sup>a</sup>	2.45 ± 0.40 <sup>b</sup>	12.15 ± 3.55 <sup>c</sup>	4.80 ± 0.42 <sup>a</sup>
SOD (SOD units/mg of protein)	7.73 ± 1.25 <sup>a</sup>	5.73 ± 0.32 <sup>b</sup>	18.50 ± 0.54 <sup>c</sup>	6.85 ± 2.30 <sup>a</sup>
PLA <sub>2</sub> (U/L.min)	52.5 ± 14.26 <sup>a</sup>	205.7 ± 62.8 <sup>b</sup>	63.4 ± 20.8 <sup>a</sup>	175.6 ± 52.5 <sup>b</sup>
Fructosamine (OD)	0.025 ± 0.003 <sup>a</sup>	0.055 ± 0.007 <sup>b</sup>	0.033 ± 0.008 <sup>a</sup>	0.63 ± 0.008 <sup>b</sup>
TBARS nmol /mg of protein	0.18 ± 0.022 <sup>a</sup>	0.30 ± 0.041 <sup>b</sup>	0.33 ± 0.042 <sup>b</sup>	0.53 ± 0.082 <sup>c</sup>

Values are mean ± SEM (n = 10 animals per group). Different letters represent significant differences (p < 0.05) for each biochemical parameter. One unit of SOD enzymatic activity is equal to the amount of enzyme that diminishes the initial absorbance of nitro blue tetrazolium by 50 %. CAT activity is expressed as the first-order constant that describes the decomposition of hydrogen peroxide at room temperature. OD, optical density; SOD, superoxide dismutase; TBARS, thiobarbituric acid-reactive substances.

### Histological study

The results of the histological study in renal tissue are shown in Table 3 and Figure 1. The different kind of lesions found were tubular tumefaction, tubular disorganization and osmotic nephrosis (lesions that precede acute tubular necrosis). In the negative control group, only 5 % moderate tubular tumefaction was found. The positive control and O<sub>2</sub> groups had similar results, with the presence of intense tubular tumefaction, osmotic nephrosis and tubular disorganization. However, in the O<sub>3</sub> group, the higher percent of lesions corresponded to a moderate tubular tumefaction, with a lower percent of osmotic nephrosis and without tubular disorganization. These results corroborate that ozone oxidative post-conditioning is able to diminish the damage caused by the I/R process.

Table 3. Results of the histological study

Groups (n=10)	MTT (%)	ITT (%)	ON (%)	TD (%)
Negative control	3	—	—	—
Positive control	—	50	40	10
O <sub>3</sub>	65	25	10	—
O <sub>2</sub>	25	45	20	5

MTT-Moderate tubular tumefaction ON-Osmotic nephrosis  
ITT- Intense tubular tumefaction TD- Tubular disorganization




Figure 1. Histology of the renal cortex using hematoxyline and eosine stain(x200). A-Positive control group, 50 % intense tubular tumefaction. B-Oxygen group, 45 % intense tubular tumefaction and 25 % moderate tubular tumefaction C-Ozone group, the higher percent of lesions corresponded to moderate tubular tumefaction and a minimum of intense tubular tumefaction D-Negative control group, moderate tubular tumefaction.

### Discussion

Oxidative stress is part of the oxygen paradox, in which the reoxygenation of ischemic kidneys generates a degree of renal injury that greatly exceeds the injury induced by ischemia alone [3,5].

It has been clearly demonstrated that ozone oxidative preconditioning has a cytoprotective effect in liver and kidneys submitted to I/R challenge, activation of the endogenous antioxidant defence system produced by the ozone treatment protecting the tissue from oxidative damage [10-12,16-20]. Beneficial effects using ozone oxidative post-conditioning have also been reported in different animal models such as diabetes or using adriamycin or cisplatin as a nephrotoxic drugs [34,36]. Rectal ozone treatment after cisplatin challenge (ozone oxidative post-conditioning) effectively prevented the decrease in renal antioxidant defence system and certainly avoided the deleterious effect of the drug [36].

Our study shows for the first time a protective effect of ozone oxidative post-conditioning in renal I/R phenomona. Repeated administration of ozone via rectal application induced a sort of cross-tolerance to free radicals released after the I/R procedure, demonstrating that renal cells have become resistant to this damage.

The mechanism of protection induced by ozone oxidative post-conditioning as well as ischaemic postconditioning [21,22,37] (which reduces myocardial infarct size by up to 50 %), are not fully understood, but the procedure has been shown to target the important mediators of lethal reperfusion injury by reducing oxidative stress, as demonstrated in this study. Other factors can also be considered, such as decreasing calcium overload, improving endothelial function, increasing adenosine levels, and attenuating the apoptosis of renal cells, among others [21,22,37]. There is evidence that ozone is able to achieve calcium homeostasis in a rat model using carbon tetrachloride as a challenge [38]. Positive effects of ozone on nitric oxide and adenosine levels and in the preservation of mitochondria in hepatic

I/R processes have also been demonstrated [39,40], as well as the control of Bax protein in renal tissue in a cisplatin model [41].

At the time of reperfusion, an increase in oxygen concentration occurs in dysfunctional mitochondria, which do not make the oxygen tetravalent reduction [6]. Also, because of neutrophil activation, there is a great release of ROS, explaining the TBARS increase (used as a measure of lipid peroxidation) in renal tissue submitted to I/R [42]. Non-significant differences were achieved between the positive control and O<sub>3</sub> groups with respect to TBARS levels. This can be explained because, in its mechanism of action, ozone produces a small and controlled oxidative stress which does not saturate the antioxidant protective mechanism, but rather triggers activation of the antioxidant enzymes [10-12,16-20,23,24,36].

An increase in TBARS was also demonstrated in the O<sub>2</sub> group. This group suffered the same lesions as the positive control group, because when oxygen levels reaching the damaged tissue increase, more ROS are released and more lipid peroxidation products are formed, but with a lower antioxidant defence system to counteract the increase in ROS. In the O<sub>3</sub> group, increases in SOD and CAT activities were demonstrated, achieving the homeostasis redox relating to protection of renal tissue after ischaemia. Levels of superoxide anion will be reduced by the activation of SOD in renal tissue, diminishing the damage to cell membranes and nephron components caused by ROS. During ischaemia, superoxide anion released can react with nitric oxide to form the peroxynitrite, a very toxic species [39]. By reducing levels of superoxide anion, less peroxynitrite will be formed and so more nitric oxide will be free to produce relaxation of the renal vascular smooth muscle, contributing to the recovery of RBF and GFR, as occurred in the O<sub>3</sub> group [2,3,43]. With regards to CAT (which converts hydrogen peroxide to water and oxygen), a significant decrease occurred in the positive control group compared with all the other groups. An increase of hydrogen peroxide, which occupies the active sites of the enzyme, has been demonstrated during I/R. Saturation of CAT therefore occurs and the enzyme cannot respond to the excess of substrate (hydrogen peroxide), being essentially inactivated. This enzymatic inactivation by excess of substrate [28,29] may explain the reduced CAT activity in the positive control group.

Phospholipase A<sub>2</sub> (PLA<sub>2</sub>) is an enzyme that splits double bonds in lipids present in membranes. It has been correlated with membrane damage by means of the release of lysophospholipids and toxic metabolites related to cell lysis in inflammatory tissues and in tissues submitted to ischaemia [18,19]. The increase in PLA<sub>2</sub> activity in the positive control and O<sub>2</sub> groups suggests that this enzyme is responsible, in combination with other factors, for the tissue damage observed in these two groups, demonstrated by histological studies. However, the enzyme activity was similar in the negative control and O<sub>3</sub> groups.

Fructosamine is an advanced glycosylation protein products highly related with protein damage and with blockade of the nitric oxide vasodilator response. Fructosamine levels were significantly higher in the positive control and O<sub>2</sub> groups, demonstrating the non-protection found in both these groups. However, fructosamine levels were similar in the O<sub>3</sub> and negative control groups.

Analyzing the results of renal function, significant decreases in RPF and GFR occurred in the O<sub>2</sub> and positive control groups, compared with the O<sub>3</sub> and negative control groups. Ischaemia damages the endothelial cells and when the blood flow decreases, shearing forces increase, producing more ROS release, damaging even more of the endothelium [39]. Ozone oxidative post-conditioning also influences tissue oxygenation via its hemorheological effects, increasing erythrocyte pliability, possibly diminishing blood viscosity and erythrocyte aggregation, allowing a higher renal blood flow during reperfusion in an organ previously submitted to a lack of blood flow [18,19].

Creatinine is not reabsorbed and is poorly secreted and so depends mainly on glomerular filtration for its excretion [1,2]. When the glomerular filtration index diminishes, creatinine excretion also decreases, leading to an increase the plasma concentration. Creatinine levels were lower in the O<sub>3</sub> group than in the positive control and O<sub>2</sub> groups, demonstrating and improvement in renal function.

Histological studies showed that lesions were most severe in the positive control and O<sub>2</sub> groups (intense tubular tumefaction, osmotic nephrosis, accompanied of tubular disorganization), correlating well with the groups that did not show a cytoprotective effect. Lesions were moderate in the O<sub>3</sub> group, proving that the renal damage is minor compared with the positive control and O<sub>2</sub> groups. From a structural point of view, ozone treatment favours kidney recovery.

### Conclusions

Ozone oxidative post-conditioning favours the functional and structural recovery of renal tissue submitted to 60 min ischaemia and 10 days reperfusion, through mechanisms that promote the maintenance of adequate cellular redox balance and improvements in blood circulation and oxygen metabolism. This is the first demonstration of a protective effect of ozone oxidative post-conditioning in renal I/R. Therefore, ozone therapy may be considered as a potential medical approach against kidney I/R damage, improving kidney transplantation. These results allow us to consider other potential medical applications for ozone therapy, mainly in the cardiovascular and central nervous system.

### References

1. Ramzi S, Vinay F, Tucker C. *Patología estructural y funcional*. New York:Mc Graw-Hill, 2000:1011-1012.
2. Brenner B M. *The Kidney*, 6<sup>th</sup> edn. Vol. 1. Philadelphia:Saunders, 2000: 1201- 1215.
3. Bonventre J. Mechanisms of ischemic renal failure. *Kidney Int* 1993; 43: 1160-1178.
4. Ichicawa I, Kiyama S, Yoshioka T. Renal antioxidant enzymes:their regulation and function. *Kidney Int* 1994; 45(1): 1- 9.
5. Annuk M, Fellstrom B, Akerblom O, et al. Oxidative stress markers in pre-uremic patients. *Clin Nephrol* 2001; 56(4): 308-314.
6. Bocci V. *Ozone. A new medical drug*. AA Dordrecht, The Netherlands:Springer Publishers, 2005:1-85.
7. Gobé G, Willgoss D, Hogg N, et al. Cell survival or death after ischemia-reperfusion injury. *Kidney Int* 1999; 56: 1299-1304.
8. Caballero M E, Calunga J L, Barber E, et al. Epidermal growth factor- mediated prevention of renal ischemia/reperfusion injury. *Biotec Ap* 2000; 17: 161-165.
9. Hammerman M R, Safirstein R, Harris R C, et al. Acute renal failure. The role of growth factors in the process of renal regeneration and repair. *Am J Physiol* 2000; 279: 3- 11.
10. Peralta C, León O S, Xaus C, et al. Protective effect of ozone treatment on the injury associated with hepatic ischemia-reperfusion: Antioxidant-Prooxidant balance. *Free Radical Res*1999; 31: 191-196.
11. Peralta C, León O S, Xaus C, et al. Effect ozone treatment on reactive oxygen species and adenosine production during hepatic ischemia-reperfusion. *Free Radical Res* 2000; 33: 595-605.
12. Ajamieh H H, Berlanga J, Merino N, et al. Role of protein synthesis in the protection conferred by ozone-oxidative-preconditioning hepatic ischaemia/reperfusion. *Transpl Int*. 2005; 18(5): 604-612.

13. Murry C E, Richard V J, Reimer K A and Jennings R B. Ischemic preconditioning slows energy metabolism and delays ultrastructural damage during a sustained ischemic episode. *Circ Res* 1990; 66: 913-931.
14. Neschis D G, Safford S D, Raghunath P N, et al. Thermal preconditioning before rat arterial balloon injury: limitation of injury and sustained reduction of intimal thickening. *Thromb Vasc Biol* 1998; 18: 120-126.
15. Riepe M W and Ludolph A C. Chemical preconditioning: a cytoprotective strategy. *Mol Cell Biochem* 1997; 174: 249-254.
16. León O S, Menéndez S, Merino N, et al. Ozone oxidative preconditioning: a protection against cellular damage by free radicals. *Mediat Inflamm* 1998; 7: 289-294.
17. Ajamieh H, Merino N, Candelario-Jalil E, et al. Similar protective effect of ischemic and ozone oxidative preconditionings in liver ischaemia/reperfusion injury. *Pharmacol Res* 2002; 45(4): 333-339.
18. Barber E, Menéndez S, León O S, et al. Prevention of renal injury after induction of ozone tolerance in rats submitted to warm ischaemia. *Mediat Inflamm* 1999; 8: 37-41.
19. Calunga J L, Menéndez S, Barber E, et al. Ozone biological response in kidneys of rats submitted to warm ischemia. *Ozone-Sci Eng* 2000; 25: 233-240.
20. Borrego A, Zamora Z, González R, et al. Protection by ozone preconditioning is mediated by antioxidant system in cisplatin induced nephrotoxicity in rats. *Mediat Inflamm* 2004; 13(1): 13-19.
21. Zhao Z Q, Corvera J S, Halkos M E, et al. Inhibition of myocardial injury by ischemic postconditioning during reperfusion: comparison with ischemic preconditioning. *Am J Physiol Heart Circ Physiol* 2003; 285: H579-H588, [Erratum, *Am J Physiol Heart Circ Physiol* 2004; 286: H477].
22. Ma X, Zhang X, Li C, Luo M. Effect of postconditioning on coronary blood flow velocity and endothelial function and LV recovery after myocardial infarction. *J Interv Cardiol* 2006; 19: 367-375.
23. Hernández F, Menéndez S and Wong R. Decrease of blood cholesterol and stimulation of antioxidative response in cardiopathy patients treated with endovenous ozone therapy. *Free Rad Biol Med* 1995; 19: 115-119.
24. Al-Dalain S M, Martínez G, Menéndez S, et al. Therapeutic efficacy of ozone medical treatments in patients with diabetic foot. *Eur J Pharmacol* 2005; 523: 151-161.
25. Bratton A C and Marshall E K. A new coupling component for sulfanilamide determination. *J Biol Chem* 1939; 128: 537-542.
26. Smith H W and Tinkelstin N. The renal clearance of substitute hippuric acid derivative and other aromatics acids in dog and man. *J Clin Invest* 1945; 24: 388-391.
27. Schreiner G. Determination of inulin by means of resorcinol. *Proc Soc Exp Biol Med* 1950; 1: 117-120.
28. Minami M and Yoshikawa H. A simplified assay method of superoxide dismutase activity for clinical use. *Clin Chim Acta* 1979; 92: 337.
29. Rice E C and Diplock A T. Laboratory techniques in biochemistry and molecular biology. In: Burtin R H, Knippenbrg P H, ed. *Techniques in free radical research*. Amsterdam:Elsevier, 1991: 199-201.
30. Ohkawa H, Orichi N and Yagi K. Assay for lipid peroxidation animal an tissues by thiobarbituric acid reaction. *Anal Biochem* 1979; 95: 351-358.
31. Hotter G, León O S and Catafau-Roselló. Tissular prostanoid release phospholipase A<sub>2</sub> activity and lipid peroxidation in pancreas transplantation. *Transplantation* 1991; 51: 987-990.
32. Lowry O H, Rocebrogh N I, Farr A L and Randall R I. Protein measurement with the folin fenol reagent. *J Biol Chem* 1951; 193: 165-175.
33. Thome J, Münch G, Müller R. Advanced glycation endproducts-associated parameters in the peripheric TAU blood of patients with Alzheimer's disease. *Life Sc* 1996; 59: 679-685.

34. Al-Dalain S M, Martínez G, Candelario-Jalil E, et al. Ozone treatment reduces markers of oxidative and endothelial damage in an experimental diabetes model in rats. *Pharmacol Res* 2001; 44(5): 391-396.
35. Calunga J L, García M, Chaple M, et al. Ozonoterapia en la glomerulonefritis tóxica experimental por adriamicina. *Rev Cubana Invest Biomed* 2004; 23(3): 139-43.
36. González R, Borrego A, Zamora Z, et al. Reversion by ozone treatment of acute nephrotoxicity induced by cisplatin in rats. *Mediat Inflamm* 2004; 13(5/6): 307-312.
37. Yellon D M and Hausenloy D J. Myocardial reperfusion injury. *N Engl Med* 2007; 357: 1121-1135.
38. León O S, Menéndez S, Merino N, et al. Influencia del preconditionamiento oxidativo con ozono sobre los niveles de calcio. *Rev CENIC Ciencias Biol* 1993; 29(3): 134-136.
39. Ajamieh H H, Menéndez S, Martínez-Sánchez G, et al. Effects of ozone oxidative preconditioning on nitric oxide generation and cellular redox balance in a rat model of hepatic ischaemia-reperfusion. *Liver Int* 2004; 24: 55-62.
40. León O S, Ajamieh H H, Berlanga J, et al. Ozone oxidative preconditioning is mediated by A<sub>1</sub> Adenosine receptors in a rat model of liver ischemia/reperfusion. *Transplant Int* 2008; 21: 39-48.
41. Borrego A, Zamora Z, González R, et al. Ozone/Oxygen mixture modifies the subcellular redistribution of Bax protein in renal tissue from rats treated with cisplatin. *Arch Med Res* 2006; 37: 717-722.
42. Paller M S. Effect of neutrophil depletion on ischemia renal injury in the rat. *J Lab Clin Med* 1993; 11(3): 379-386.
43. Rodríguez-Puyol D, Lucio J, Ruiz P, et al. Radicales libres y daño glomerular. *Nefrología* 1996; XVI(S3): 29-34.